



High-cadence observations of galactic nuclei by the future two-band UV-photometry mission *QUVIK*



Michal Zajaček¹, Norbert Werner¹, Aaron Tohuvavohu² et al.
(on behalf of *QUVIK* Science Team)

¹Masaryk University, ²University of Toronto

May 9, 2024 - UV Science and Instrumentation Workshop, NASA JPL

QUVIK – Quick Ultra-Violet Kilonova surveyor

- **QUVIK** was approved for funding by the Czech Ministry of Transport and the European Space Agency as an **ambitious Czech national space mission**
- **primary objective:** **kilonovae** following the mergers of neutron stars
- **secondary objectives:** hot stars, supernovae, star clusters, γ -ray bursts, and **(active) galactic nuclei**
- modified Cassegrain of ~ 33 cm; $\text{FOV} \sim 1^\circ \times 1^\circ$, $\text{PSF} \lesssim 2.5''$
- two bands: **NUV** (260 - 360 nm) and **FUV** (140 - 190 nm)



QUVIK – Quick Ultra-Violet Kilonova surveyor

- scientific targets and program elaborated in three review papers published in the **Space Science Reviews** special issue on **QUVIK**

1. mission description and stellar transients (2306.15080),
2. observations of stars and stellar systems (2306.15081),
3. active galactic nuclei and nuclear transients (2306.15082)

Space Science Reviews (2024) 205:11
https://doi.org/10.1007/s11214-024-01008-1

Science with a Small Two-Band UV-Photometry Mission I:
Mission Description and Follow-up Observations of Stellar
Transients

N. Werner¹ · J. Ripa² · C. Thöne³ · F. Mätz⁴ · R. Karfirt⁵ · M. Jeliński⁶ · F. Hoesch⁷ ·
J. Bendažek⁸ · M. Topinka⁹ · G. Lukes-Gerolopoulos¹⁰ · M. Zajaček¹¹ · M. Lohaj¹² ·
M. Pršegan¹³ · A. Růžička¹⁴ · J. Mész¹⁵ · A. Růž¹⁶ · O. Popely¹⁷ · A. Banaś¹⁸ · J. Jan¹⁹ ·
R. Seidemann²⁰ · J. Geronzi²¹ · A. Vlahakis²² · L. Stojan²³ · J. Segura²⁴ · E. Bahar²⁵ ·
S. Tavec²⁶ · J. Loh²⁷ · O. Hach²⁸ · S. Ben Ami²⁹ · M. J. Baruch³⁰ · O. Berger³¹ ·
A. Tatarski³² · S. Suvandani³³ · M. Bala³⁴ · M. H. S. Papp³⁵ ·
Huanyang Kuang Chang³⁶

Received: 27 June 2023 / Accepted: 9 January 2024 / Published online: 2 February 2024
© The Author(s) 2024

Abstract

This is the first in a collection of three papers introducing the science with an ultra-violet (UV) space telescope on an approximately 100 kg small satellite with a moderately fast re-pointing capability and a real-time alert communication system approved for a Czech national space mission. The mission, called Quick Ultra-Violet Kilonova surveyor – QUVIK, will provide low follow-up capabilities to increase the discovery potential of gravitational wave observations and future wide-field multi-wavelength surveys. The primary objective of the mission is the measurement of the UV brightness evolution of kilonovae, resulting from mergers of neutron stars, to distinguish between different ejection scenarios. The mission, which is designed to be complementary to the *Ultaviolet Transient Astronomy Satellite* – *UTASAT*, will also provide follow-up capabilities for other transients both in the near- and far-UV bands. Between the observations of transients, the satellite will target other objects described in this collection of papers, which demonstrates that a small and relatively affordable orbital UV-space telescope can be transformative for many fields of astrophysics.

Keywords UV space observatory · Kilonovae · Gamma-ray bursts · Supermassive

1 Introduction

The first simultaneous detection of gravitational waves and electromagnetic radiation on 2017 August 17 (Abbott et al. 2017a), resulting from a coalescence of neutron stars, marked the onset of multi-messenger astrophysics involving gravitational waves. This exciting observation showed that neutron stars mergers are of major importance for studying the Universe with heavy elements created as gold and platinum. The radioactive decay of these heavy elements powers a thermal outburst at ultra-violet/visible/near-infrared wavelengths

Extended author information available on the last page of the article.

Space Science Reviews (2024) 205:29
https://doi.org/10.1007/s11214-024-01008-1

Science with a Small Two-Band UV-Photometry Mission II:
Observations of Stars and Stellar Systems

J.H. Kriváček¹ · Jan Bendažek^{2,3} · Jan Budaj⁴ · Daniela Kostalová⁵ · Andrius Pliūš⁶ ·
Martín Peckol⁷ · Miloslav Zdražil⁸ · Václav Bakoš⁹ · Miroslav Bradač¹⁰ ·
Huanyang Kuang Chang¹¹ · Nida Fabbro¹² · Rastislav Gálik¹³ · Daniel Jacholkowski¹⁴ ·
Jan Jaroš¹⁵ · Jan Kral¹⁶ · Jakub Kral¹⁷ · Jan Kratoch¹⁸ ·
Brenkita Khabibulaeva¹⁹ · Peter Kufner²⁰ · Miroslav Labaj²¹ · Jovana Mész²² ·
Zdeněk Mikulášek²³ · Filip Mlýns²⁴ · Joss Fausnaugh²⁵ · Michal Pršegan²⁶ ·
Tahereh Ramezani²⁷ · Terence Regierová²⁸ · Jakub Ripa²⁹ · Linda Schmidtke-Buchatz³⁰ ·
Mark Skaržek³¹ · Gabriel Szász³² · Werner Wimmer³³ · Michal Zajaček³⁴ · Norbert Werner³⁵

Received: 26 June 2023 / Accepted: 23 February 2024 / Published online: 13 March 2024
© The Author(s) 2024

Abstract

We outline the impact of a small two-band UV-photometry satellite mission on the field of stellar physics, spectroscopy of stars, binaries, stellar clusters, interstellar matter, and exoplanets. On specific examples of different types of stars and stellar systems, we discuss particular requirements for such a satellite mission in terms of specific mission parameters, such as bandwidth, precision, cadence, and observation duration. We show that such a mission may provide essential data not only for hot stars but also most of their light in UV, but also for cool stars, where UV traces their activity. This is important, for instance, for exoplanet studies, because the level of stellar activity influences habitability. While the main asset of the two-band UV mission lies in time-domain astronomy, an example of open clusters proves that such a mission would be important also for the study of stellar populations. Properties of the interstellar dust are best explored when combining optical and IR information with observations in UV.

It is well known that data about UV radiation evolution. Consequently, we outline how such a UV mission can be used to detect objects of sufficiently hot stars by various data objects and study disks, rings, clumps, disintegrating exoplanets or comets. Furthermore, UV radiation can be used to study the cooling of neutron stars providing information about the extreme states of matter in the interiors of neutron stars and used for mapping heated spots on their surfaces.

Keywords Techniques: photometric · Chromospheres · Stars: variables: general · Binaries: general · Open clusters and associations: general · Planetary systems

1 Introduction

The sun discloses its astrophysical changes during the last few decades more frequently connected with the opening of new observational windows into invisible parts of the spectrum. Recently, the advent of observations working during the electromagnetic domain founded a

Extended author information available on the last page of the article.

Space Science Reviews (2024) 205:29
https://doi.org/10.1007/s11214-024-01008-1

Science with a Small Two-Band UV-Photometry Mission III:
Active Galactic Nuclei and Nuclear Transients

M. Zajaček¹ · B. Corry² · K.K. Jahnke³ · M. Stach^{4,5} · U. Koss⁶ · A. Pender⁷ ·
D.R. Pasham⁸ · M. Srinivasan⁹ · V. Witzany¹⁰ · P. Suková¹¹ · F. Mész¹² · N. Wimmer¹³ ·
J. Ripa¹⁴ · J. Mész¹⁵ · M. Labaj¹⁶ · P. Karfirt¹⁷ · J. Kriváček¹⁸

Received: 20 June 2023 / Accepted: 13 March 2024 / Published online: 28 March 2024
© The Author(s) 2024

Abstract

In this article, the third one in the series focused on a small two-band UV-photometry mission, we discuss possibilities for a small UV two-band photometry mission in studying accreting supermassive black holes (SMBHs, mass range $\sim 10^6$ – $10^9 M_\odot$). We focus on the following observational concepts: (i) detection and monitoring of sub-keV type I Active Galactic Nuclei (AGN) in order to measure the time delay between the far-UV, the near-UV, and other wavelengths (UV-opt and optical); (ii) monitor transients including spatially rapid disruption events and repetitive nuclear transients; and (iii) the study of peculiar sources, such as changing-look AGN, blazars and gaps in accretion disks, low-luminosity AGN, and candidates for Intermediate-Mass Black Holes (IMBHs, mass range $\sim 10^3$ – $10^4 M_\odot$) in galactic nuclei. The importance of a small UV mission for the observing program (ii) is to provide intense, high-cadence monitoring of selected sources, which will be beneficial for, e.g., reverberation-mapping of accretion disks and subsequently constraining accretion disk models with observations. For program (ii), a relatively small UV space telescope is suitable even for such monitoring a transient event within $\lesssim 20$ minutes after receiving the trigger, such a moderately fast re-pointing capability will be highly beneficial. Peculiar sources within the program (iii) will be of interest to a wider community and will create an environment for competitive observing proposals. For total disruption events (TDEs), high-cadence UV monitoring is crucial for distinguishing among different scenarios for the origin of the UV emission. The small two-band UV space telescope will also provide information about the near- and far-UV continuum variability for rare transients, such as active galactic TDEs and jetted TDEs. We also discuss the possibilities to study and analyze sources with non-steady accretion flows, such as AGN with puppy disks, low-luminosity active galactic nuclei with intermittent accretion, and SMBH binaries potentially involving intermediate-mass black holes.

Keywords Galactic nuclei · Accretion flows · Tidal disruption events · Transients · Photometry · Time series


1 Introduction

The growth of supermassive black holes (hereafter SMBHs) residing in the centers of galaxies is a crucial topic in modern astrophysics (D. Martin 2019). SMBHs can grow by accre-

Extended author information available on the last page of the article.


QUVIK – Quick Ultra-Violet Kilonova surveyor

For technical details concerning the *QUVIK* NUV and FUV cameras, see the **poster “NUV and FUV cameras for the Quick Ultra-Violet Kilonova Surveyor (QUVIK)”** by **Aaron Tohuavohu** (University of Toronto)





QUVIK Galactic nuclei program

- **monitoring of selected AGN:** photometric reverberation mapping
- **nuclear transients:** tidal disruption events - TDEs, changing-look AGN, and peculiar sources



QUVik Galactic nuclei program


- accretion disk UV emission (type I AGN) is observable up to $z \sim 0.5$
- depends on the SMBH mass ($10^7 - 10^8 M_\odot$) and the relative accretion rate (0.1 – 1.0)



Redshift limit	Limiting u magnitude	Number of SDSS quasars	$t_{\text{int}} [\text{s}], S/N = 100$
0.5	17.0	151	100.0
0.5	18.0	964	251.2
0.7	17.0	167	100.0
0.7	18.0	1047	251.2

Dedicated monitoring

- **Spatial resolution** $\Leftrightarrow$ **Temporal resolution**
- Different wavelengths probe different scales of an accretion disk and its immediate surroundings



Dedicated monitoring

- driving ionizing radiation ΔF_i (X-ray corona)
- reprocessed radiation in the far-UV, near-UV, optical domains ΔF_r

■


$$\Delta F_r(t) = \int_0^{\tau_{\max}} \psi(\tau) \Delta F_i(t - \tau) d\tau,$$

where $\tau = r/c$ is the mean time-delay due to light-travel time, $\psi(\tau)$ is the transfer function of the accretion disk (delay and blurring)

Reverberation mapping		Light-crossing time [days]		
Wavelength domain	Spatial length scale [r_g]	$10^7 M_\odot$	$10^8 M_\odot$	$10^9 M_\odot$
X-ray	1–10	5.7×10^{-4} – 5.7×10^{-3}	5.7×10^{-3} – 5.7×10^{-2}	5.7×10^{-2} –0.57
UV/optical (QUVik)	10^2 – 10^4	5.7×10^{-2} –5.7	0.57–57	5.7–570
optical BLR	10^3 – 10^5	0.57–57	5.7–570	57–5700
optical/infrared dusty torus	$> 10^5$	> 57	> 570	> 5700

Dedicated monitoring

- mock observation with different cadences and SMBH masses of $10^7 M_{\odot}$ and $10^8 M_{\odot}$
- light curve of the X-ray corona calculated using the Timmer-König algorithm





Dedicated monitoring

- for $M_{\bullet} = 10^7 M_{\odot}$, 10-day monitoring with a high cadence of 0.1 days is sufficient
- for $M_{\bullet} = 10^8 M_{\odot}$, cadence can be lower (~ 1 day), but the observation should last for about half a year

$M_{\bullet}$ ($M_{\odot}$)	S/N	RMS (%)	ΔT (days)	T (days)	τ_{ICCF} (days)	$\Delta \tau_{ICCF}$ (days)	τ_{χ^2} (days)	$\Delta \tau_{\chi^2}$ (days)	τ_{ψ} (days)
10^7	∞	11.40	0.1	10	0.172	0.086	0.135	0.047	0.205
10^7	100	11.50	0.1	10	0.165	0.069	0.125	0.072	0.205
10^8	∞	3.15	0.25	186	1.021	0.182	0.686	0.056	0.786
10^8	100	3.51	0.25	186	0.966	0.193	0.621	0.001	0.786
10^8	∞	3.15	0.5	186	1.028	0.183	0.708	0.047	0.786
10^8	100	3.33	0.5	186	1.017	0.278	1.112	0.250	0.786
10^8	∞	3.15	1.0	186	1.087	0.218	0.698	0.045	0.786
10^8	100	3.32	1.0	186	1.065	0.319	0.527	0.049	0.786

Dedicated monitoring

- potential effect of extended media (UFO, BLR) – prolongation with respect to the time delay due to disk reprocessing,
 $\tau(\lambda) \propto \lambda^{4/3}$
- redshift-dependent, needs further numerical assessment




- calculations based on the **Disk+BLR transfer function** (left) and obtained numerically using **stochastic variability** (right)


TDE/nuclear transients

- tidal disruption events are accompanied by bright optical/UV emission that decreases as a power-law ($\dot{M}_{\text{fallback}} \propto t^{-5/3}$)
- **unclear origin of the UV emission:** shock-produced or accretion-disk emission?


Shock at apocenter




Series of discrete self-interactions



Elliptical disc




Reprocessing scenario





TDE/nuclear transients

- high-cadence monitoring shortly after the optical/UV outburst could reveal **quasiperiodic modulations due to the Lense-Thirring precession $\Rightarrow$ constraining SMBH spin**
- for a given periodicity, spin constraints depend on the post-TDE disk parameters (density slope, outer radius) and the SMBH mass

(a) Lower luminosity and temperature




(b) Higher luminosity and temperature



Non-standard accretion flows: low-luminosity AGN



- constraining SEDs of nearby low-luminosity AGN ($10^{-3} \lesssim \dot{m} \lesssim 10^{-2}$)
- contribution of starlight is more relevant and eventually may prevent detecting a fainter AGN



M81 is a nearby (12 Mly) grand spiral with a small jet: suitable source for the disentangling of the host from the core (image credit: HST, GALEX, Spitzer)


Non-standard accretion flows: perturbation by an orbiting IMBH

- for an AGN disk, an orbiting IMBH ($m_{\text{IMBH}} \gtrsim 10^3 M_{\odot}$) that passes through the disk can produce **quasiperiodic UV flashes** (see also Linial & Metzger 2024 for the analysis of UV Quasiperiodic Erupters - QPEs)
- function of an IMBH mass and its inclination



Summary

- **reverberation mapping** of the accretion disk between FUV and NUV bands (and other wavebands)
- origin of the **TDE *UV emission***
- **non-standard accretion**: low-luminosity AGN, changing-look AGN, binary supermassive black holes (“gappy” disks)
- **unexpected discoveries**, e.g. quasiperiodic transients with UV flashes due to an orbiting and inclined intermediate-mass black hole / UV Quasiperiodic Eruptors



Credit: R. Hurt (Caltech/IPAC)

**MASARYK
UNIVERSITY**